

NIKIWE WOMEN'S SAFE ABORTION CLINIC IN CAPE TOWN ((0735521667

DR NIKIWE

Tel: 0735521667



Location

Western Cape, Cape Town

<https://www.freeadsz.co.za/x-340890-z>

NIKIWE WOMEN'S SAFE ABORTION CLINIC /TERMINATION OF PREGNANCY 0735521667
SAME DAY PAIN FREE

NIKIWE Women's Safe Abortion Clinic / Termination of pregnancy Clinic/ Abortion pills .WE ALSO
TELEMEDICINE WITH DOOR TO DOOR AND POSTAL DELIVERY

Medical Abortion and surgical

What is a medical abortion?

- A medical abortion uses 2 different medications, The medications end a pregnancy and make the uterus push out the pregnancy tissue (like a miscarriage).

Can I have a medical abortion?

- You may be able to have a medical abortion if:

- › Your pregnancy is 2 weeks- 22 weeks counting from the first day of your last regular period.

- › Your pregnancy is in your uterus, not in your fallopian tubes (ectopic pregnancy).

- › You do not have an intrauterine device (IUD) in place.

- When you meet with a health care provider, they will ask you about your health

and any prescription medications you are taking. This will help them find out if you are eligible for a medical abortion.

Important

Medical abortion can cause very strong pain and bleeding. Make sure that you can manage these symptoms at home. You may need to come to a procedural sedation instead. Any concerns with your health care provider will help decide.


In rare cases, you may have very heavy bleeding that needs medical attention. For this reason, if you have a medical abortion, you must be able to get to an Emergency Department right away.

What happens during a medical abortion?

- You will have an appointment with a health care provider to talk about the




NIKIWE WOMEN'S
SAFE ABORTION CLINIC IN
CAPE TOWN ((0735521667
DR NIKIWE)) ; SAME
<https://www.freeadsz.co.za/x-340890-z>



NIKIWE WOMEN'S
SAFE ABORTION CLINIC IN
CAPE TOWN ((0735521667
DR NIKIWE)) ; SAME
<https://www.freeadsz.co.za/x-340890-z>



NIKIWE WOMEN'S
SAFE ABORTION CLINIC IN
CAPE TOWN ((0735521667
DR NIKIWE)) ; SAME
<https://www.freeadsz.co.za/x-340890-z>



NIKIWE WOMEN'S
SAFE ABORTION CLINIC IN
CAPE TOWN ((0735521667
DR NIKIWE)) ; SAME
<https://www.freeadsz.co.za/x-340890-z>



NIKIWE WOMEN'S
SAFE ABORTION CLINIC IN
CAPE TOWN ((0735521667
DR NIKIWE)) ; SAME
<https://www.freeadsz.co.za/x-340890-z>

procedure in detail.

2

- You will be given a prescription for a medication called Mifegymiso.
 - › This prescription can be filled at any pharmacy.

What supplies will I need before taking Mifegymiso?

- You will need several large period (maxi) pads. Do not use tampons or period cups during your medical abortion.
- You may take over-the-counter pain medication like acetaminophen (Tylenol®), ibuprofen (Advil ®, Midol®), or naproxen (Aleve®). Be sure to have these medications on hand before starting.
- Other comfort measures, like a heating pad or a hot water bottle, can also help with pain.

How to take the medications

Step 1: Mifepristone (green box)

- Swallow 1 mifepristone tablet with water.
- Once you take mifepristone, there is no way to reverse the effects. This tablet will end the pregnancy.
- Most people do not feel different after taking this tablet. A small number of people have mild bleeding or cramping.

After taking mifepristone,

Step 2 : Misoprostol (orange box)

- Misoprostol softens the cervix and tells your body to push out the tissue (like a miscarriage). You will have, painful cramps during this step.
- Take pain medication about 30 minutes before taking misoprostol:
 - › For example, you can take both acetaminophen (1000 mg) and ibuprofen (400 to 800 mg) with food.
 - There are 2 ways to take misoprostol. If you have nausea (feeling sick to your stomach), it is recommended to take misoprostol vaginally (option 1). Both ways work equally well.

3

Option 1. Vaginally

After you take the mifepristone tablet
insert (put) all 4 misoprostol tablets in your vagina. ***** them as high up as you can, like you ***** a tampon or a period cup.

Option 2. Orally (by mouth)

After you take the mifepristone tablet (Step 1 on page 2), place all 4 misoprostol tablets in your mouth at the same time. Place 2 tablets between your cheek and gums on each side. Do not swallow these tablets. Leave them

there for 30 minutes. After 30 minutes, drink some water to swallow any pieces that are left.

What will happen after taking misoprostol?

- Within 1 to 3 hours of taking misoprostol, you will start to have normal cramping, and bleeding, and pass clots and tissue. This is the tissue starting to empty from the uterus.
- You may have very heavy bleeding with clots as small as quarters,
- You may have bleeding and cramping for a few hours (30minutes -3hours), until all the tissue passes.

Side effects

- After taking misoprostol, you may have:
 - › A fever
 - › Chills
 - › Diarrhea (loose, watery poop)

These symptoms are common. If you are worried about any of these symptoms, call or text the Nikiwe Clinic

Follow-up appointment(s)

- You will have a follow-up appointment 7 to 14 days after you take mifepristone (the first medication in the green box).
 - › During this appointment, the doctor will check that your medical abortion worked. You may need another blood test.
 - › After your blood test, a nurse will call to confirm whether your medical abortion is complete and answer any questions you may have.

4

What are the possible risks?

- There is a 0.5% chance that the medication will not work and you will still be pregnant. If this happens, it is recommended that you have more pills