


# NIKIWE WOMEN&apos;S CLINIC SAFE ABORTION CLINIC IN CARLETONVILLE +2773

DR.NIKIWE  
Tel: +27735521667



Location **North-West, Carletonville**  
https://www.freeadsz.co.za/x-341848-z

NIKIWE WOMEN&apos;S SAFE ABORTION CLINIC /TERMINATION OF PREGNANCY 0735521667  
SAME DAY PAIN FREE

NIKIWE Women"s Safe Abortion Clinic / Termination of pregnancy Clinic/ Abortion pills .WE ALSO  
TELEMEDICINE WITH DOOR TO DOOR AND POSTAL DELIVERY

Medical Abortion and surgical











What is a medical abortion?

- A medical abortion uses 2 different medications, The medications end a pregnancy and make the uterus push out the pregnancy tissue (like a miscarriage).

Can I have a medical abortion?

- You may be able to have a medical abortion if:
  - › Your pregnancy is 2 weeks- 22 weeks counting from the first day of your last regular period.
  - › Your pregnancy is in your uterus, not in your fallopian tubes (ectopic pregnancy).
  - › You do not have an intrauterine device (IUD) in place.
- When you meet with a health care provider, they will ask you about your health

and any prescription medications you are taking. This will help them find out if you are eligible for a medical abortion.

 <p>NIKIWE WOMEN&amp;apos;S CLINIC SAFE ABORTION CLINIC IN CARLETONVILLE DR.NIKIWE 667 SAME DAY https://www.freeadsz.co.za/x-341848-z</p>	 <p>NIKIWE WOMEN&amp;apos;S CLINIC SAFE ABORTION CLINIC IN CARLETONVILLE DR.NIKIWE 667 SAME DAY https://www.freeadsz.co.za/x-341848-z</p>	 <p>NIKIWE WOMEN&amp;apos;S CLINIC SAFE ABORTION CLINIC IN CARLETONVILLE DR.NIKIWE 667 SAME DAY https://www.freeadsz.co.za/x-341848-z</p>	 <p>NIKIWE WOMEN&amp;apos;S CLINIC SAFE ABORTION CLINIC IN CARLETONVILLE DR.NIKIWE 667 SAME DAY https://www.freeadsz.co.za/x-341848-z</p>	<p>Important</p>  <p>NIKIWE WOMEN&amp;apos;S CLINIC SAFE ABORTION CLINIC IN CARLETONVILLE DR.NIKIWE 667 SAME DAY https://www.freeadsz.co.za/x-341848-z</p>	 <p>NIKIWE WOMEN&amp;apos;S CLINIC SAFE ABORTION CLINIC IN CARLETONVILLE DR.NIKIWE 667 SAME DAY https://www.freeadsz.co.za/x-341848-z</p>	 <p>NIKIWE WOMEN&amp;apos;S CLINIC SAFE ABORTION CLINIC IN CARLETONVILLE DR.NIKIWE 667 SAME DAY https://www.freeadsz.co.za/x-341848-z</p>	 <p>NIKIWE WOMEN&amp;apos;S CLINIC SAFE ABORTION CLINIC IN CARLETONVILLE DR.NIKIWE 667 SAME DAY https://www.freeadsz.co.za/x-341848-z</p>	 <p>NIKIWE WOMEN&amp;apos;S CLINIC SAFE ABORTION CLINIC IN CARLETONVILLE DR.NIKIWE 667 SAME DAY https://www.freeadsz.co.za/x-341848-z</p>	 <p>NIKIWE WOMEN&amp;apos;S CLINIC SAFE ABORTION CLINIC IN CARLETONVILLE DR.NIKIWE 667 SAME DAY https://www.freeadsz.co.za/x-341848-z</p>
---	--	--	--	--	--	--	--	--	--

---

procedure in detail.

2

- You will be given a prescription for a medication called Mifegymiso.
- › This prescription can be filled at any pharmacy.

What supplies will I need before taking Mifegymiso?

- You will need several large period (maxi) pads. Do not use tampons or period cups during your medical abortion.
- You may take over-the-counter pain medication like acetaminophen (Tylenol®), ibuprofen (Advil®, Midol®), or naproxen (Aleve®). Be sure to have these medications on hand before starting.
- Other comfort measures, like a heating pad or a hot water bottle, can also help with pain.

How to take the medications

Step 1: Mifepristone (green box)

- Swallow 1 mifepristone tablet with water.
- Once you take mifepristone, there is no way to reverse the effects. This tablet will end the pregnancy.
- Most people do not feel different after taking this tablet. A small number of people have mild bleeding or cramping.

After taking mifepristone,

Step 2 : Misoprostol (orange box)

- Misoprostol softens the cervix and tells your body to push out the tissue (like a miscarriage). You will have, painful cramps during this step.
- Take pain medication about 30 minutes before taking misoprostol:
  - › For example, you can take both acetaminophen (1000 mg) and ibuprofen (400 to 800 mg) with food.
- There are 2 ways to take misoprostol. If you have nausea (feeling sick to your stomach), it is recommended to take misoprostol vaginally (option 1). Both ways work equally well.

3

Option 1. Vaginally

After you take the mifepristone tablet insert (put) all 4 misoprostol tablets in your vagina. \*\*\*\*\* them as high up as you can, like you \*\*\*\*\* a tampon or a period cup.

Option 2. Orally (by mouth)

After you take the mifepristone tablet (Step 1 on page 2), place all 4 misoprostol tablets in your mouth at the same time. Place 2 tablets between your cheek and gums on each side. Do not swallow these tablets. Leave them

---

there for 30 minutes. After 30 minutes, drink some water to swallow any pieces that are left.

What will happen after taking misoprostol?

- Within 1 to 3 hours of taking misoprostol, you will start to have normal cramping, and bleeding, and pass clots and tissue. This is the tissue starting to empty from the uterus.
- You may have very heavy bleeding with clots as small as quarters,
- You may have bleeding and cramping for a few hours (30minutes -3hours), until all the tissue passes.

Side effects

- After taking misoprostol, you may have:
  - › A fever
  - › Chills
  - › Diarrhea (loose, watery poop)

These symptoms are common. If you are worried about any of these symptoms, call or text the Nikiwe Clinic

Follow-up appointment(s)

- You will have a follow-up appointment 7 to 14 days after you take mifepristone (the first medication in the green box).
  - › During this appointment, the doctor will check that your medical abortion worked. You may need another blood test.
  - › After your blood test, a nurse will call to confirm whether your medical abortion is complete and answer any questions you may have.

4

What are the possible risks?

- There is a 0.5% chance that the medication will not work and you will still be pregnant. If this happens, it is recommended that you have more pills